In the function above, b is a constant. If f(6) = 7, what is the value of f(-2)?

- A) -5
- B) -2
- C) 1
- D) 7

4

If f(x) = -2x + 5, what is f(-3x) equal to?

- A) -6x 5
- B) 6x + 5
- C) 6x 5
- D)  $6x^2 15x$

5

$$\sqrt{2k^2 + 17} - x = 0$$

If k > 0 and x = 7 in the equation above, what is the value of k?

- A) 2
- B) 3
- C) 4
- D) 5

5

$$(x^2y - 3y^2 + 5xy^2) - (-x^2y + 3xy^2 - 3y^2)$$

Which of the following is equivalent to the expression above?

- A)  $4x^2y^2$
- B)  $8xy^2 6y^2$
- C)  $2x^2y + 2xy^2$
- D)  $2x^2y + 8xy^2 6y^2$

6

$$3x^2 - 5x + 2$$

$$5x^2 - 2x - 6$$

Which of the following is the sum of the two polynomials shown above?

- A)  $8x^2 7x 4$
- B)  $8x^2 + 7x 4$
- (C)  $8x^4 7x^2 4$
- D)  $8x^4 + 7x^2 4$

10

A function f satisfies f(2) = 3 and f(3) = 5. A function g satisfies g(3) = 2 and g(5) = 6. What is the value of f(g(3))?

- A) 2
- (B) 3
- C) 5
- D) 6

13

$$h = -16t^2 + vt + k$$

The equation above gives the height h, in feet, of a ball t seconds after it is thrown straight up with an initial speed of  $\nu$  feet per second from a height of k feet. Which of the following gives  $\nu$  in terms of h, t, and k?

- A) v = h + k 16t
- B)  $v = \frac{h k + 16}{t_{\odot}}$
- $C) \quad v = \frac{h+k}{t} 16t$
- $D) \quad v = \frac{h-k}{t} + 16t$

4

| n    | 1  | 2 | 3 | 4 |
|------|----|---|---|---|
| f(n) | -2 | 1 | 4 | 7 |

The table above shows some values of the linear function f. Which of the following defines f?

- A) f(n) = n 3
- B) f(n) = 2n 4
- C) f(n) = 3n 5
- D) f(n) = 4n 6

2 =

| x | f(x) |  |  |  |
|---|------|--|--|--|
| 0 | -2   |  |  |  |
| 2 | 4    |  |  |  |
| 6 | 16   |  |  |  |

Some values of the linear function f are shown in the table above. What is the value of f(3)?

- A) (
- B)
- CI = 8
- D) 9